ΒΙΒΛΙΟΓΡΑΦΙΑ ΑΡΘΡΩΝ

- 1. Amato L, Minozzi S, Davoli M. Efficacy and safety of pharmacological interventions for the treatment of the Alcohol Withdrawal Syndrome (Review). The Cochrane Collaboration, 2011
- 2. Babor TF & Higgins-Biddle JC. Brief Intervention for Hazardous and Harmful Drinking. A Manual for Use in Primary Care. Geneva, 2001, World Health Organization
- 3. Barber JG, Gilbertson R. An experimental study of brief unilateral intervention for the partners of heavy drinkers. Res Soc Work Pract 1996;6(3):325-36
- Berks J & McCormick R. Screening for alcohol misuse in elderly primary care patients: a systematic literature review. Int Psychogeriatr 2008; 20:6, 1090-1103
- 5. Berner MM et al. Detection and management of alcohol use disorders in German primary care influenced by non-clinical factors. Alcohol & Alcoholism, 2008; 42(4):308-316
- 6. Bradley KA, Boyd-Wickizer J, Powell SH, Burman ML. Alcohol screening questionnaires in women: a critical review. JAMA 1998;280(2):166-71
- Garbutt JC, West SL, Carey TS, Lohr KN, Crews FT. Pharmacological treatment of alcohol dependence: a review of the evidence. JAMA 1999;281(14):1318-25
- 8. Hilton ME, Maisto SA, Conigliaro J, McNiel M, Kraemer K, Kelley ME, et al. Improving alcoholism treatment across the spectrum of services. Alcohol Clin Exp Res 2001;25(1):128-35.
- 9. Kahan M, Wilson L, Becker L. Effectiveness of physician-based interventions with problem drinkers: a review. CMAJ 1995;152(6):851-9
- 10. Kaner E, Lock C, Heather N, McNamee P, Bond S. Promoting brief alcohol intervention by nurses in primary care: a cluster randomised controlled trial. Patient Educ Couns. In press 2003
- 11. Kaner E et al. Effectiveness of screening and brief alcohol intervention in primary care (SIPS trial): pragmatic cluster randomised controlled trial. BMJ 2013;346:e8501
- 12. Kypri K et al. Randomized controlled trial of web-based alcohol screening and brief intervention in primary care. Arch Intern Med 2008; 168(5):530-536
- 13. Martinus T, Anderson B, Carter H. Counseling for alcohol problems in primary care in Forth Valley an innovative approach? Health Bull (Edinb) 2001;59(3):158-62.
- 14. Mayo-Smith MF. Pharmacological management of alcohol withdrawal. A meta-analysis and evidence-based practice guideline. American Society of Addiction Medicine Working Group on Pharmacological Management of Alcohol Withdrawal. JAMA 1997;278(2):144-51
- 15. Moyer A, Finney JW, Swearingen CE, Vergun P. Brief interventions for alcohol problems: a meta-analytic review of controlled investigations in treatmentseeking and non-treatment-seeking populations. Addiction 2002;97(3):279-92

- 16. Navarro HJ et al. The cost-effectiveness of tailored, postal feedback on general practitioners' prescribing of pharmacotherapies for alcohol dependence. Drug Alcohol Depend 2012; 124:207-215
- 17. Randall CL, Thomas S, Thevos AK. Concurrent alcoholism and social anxiety disorder: a first step toward developing effective treatments. Alcohol Clin Exp Res 2001;25(2):210-20.
- 18. Salaspuro M. Carbohydrate-deficient transferrin as compared to other markers of alcoholism: a systematic review. Alcohol 1999;19(3):261-71.
- 19. Schaus JF et al. Alcohol screening and brief intervention in a college student health center: a randomized controlled trial. J Stud.Alcohol Drugs 2009; supp. 16: 131-141
- 20. Scouller K, Conigrave KM, Macaskill P, Irwig L, Whitfield JB. Should we use carbohydrate-deficient transferrin instead of gamma-glutamyltransferase for detecting problem drinkers? A systematic review and metaanalysis. Clin Chem 2000;46(12):1894-902.
- 21. Sellers EM, Naranjo CA, Harrison M, Devenyi P, Roach C, Sykora K. Diazepam loading: simplified treatment of alcohol withdrawal. Clin Pharmacol Ther 1983;34(6):822-6
- Solberg LI, Maciosek MV, Edwards NM. Primary care intervention to reduce alcohol misuse. Ranking its health impact and cost effectiveness. Am. J. Prev. Med. 2008; 34, 143–152
- 23. Stout RL, Rubin A, Zwick W, Zywiak W, Bellino L. Optimizing the cost effectiveness of alcohol treatment: a rationale for extended case monitoring. Addict Behav 1999;24(1):17-35.
- 24. Sullivan LE et al. A meta-analysis of the efficacy of nonphysician brief interventions for unhealthy alcohol use: implications for the patient-centered medical home. Am J Addict 2011; 20: 343–356
- 25. Toft U et al. The impact of a population-based multi-factorial lifestyle intervention on alcohol intake. The Inter99 study. Prev Med 2009; 115-121